440000
460000
480000
500000
6160000
6140000
6120000
6100000
6080000
6060000
Oksbol
Varde
Hjerting
Saedden
Esbjerg
Bramming
Nordby
Fano
Sonderho
Mando
Mando By
Ribe
Romo
Kongsmark
Skaerbaek
Bredebro
Logumkloster
Hojer
Tonder
Sylt
Westerland
Foehr
Abiotic features
low intertidal flats (<50% E.)
high intertidal flats (>50% E.)
sl sh sand (<5% clay)
xl xh muddy sand (5-8% clay)
ml mh sand (>8% clay)
67% emersion time
0 beach above Mean High Tide
N depth in m below Sea Level < 10
N10 depth in m below Sea Level > 10
Biotic features
musselbed
Zostera marina bed (with 200 meter buffer)
Zostera noltii (with 200 meter buffer)
1 mud- or sand flat with salt-marsh pioneers
2 low salt marsh (Puccinellion maritimae)
3 middle salt marsh (Armerion maritimae)
4 high marsh (Lolio-Potentillion anserinae)
5 transition between salt-marsh and sand dunes
6 reed marshes
HABITATS OF THE NETHERLANDS, GERMAN AND DANISH WADDEN SEA
Compiled by: K. S. Dijkema, Rijksinstituut voor Natuurbeheer, Texel
1 : 100.000
sheet 5
ALTERRA WAGENINGEN UR
nature-consult
N
GIS-Compilation:
H.-P. Dauck
WGS_1984_UTM_Zone_32N
0 1 2 3 4 5
Kilometer
nature-consult
Hackelbrink 21
D-31139 Hildesheim
www.nature-consult.de